

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

to the withdrawal of water. Gorke goes a step further and shows³⁰ that the low temperatures besides producing various analogous changes, lead to intramolecular transpositions in the proteids, which are generally accompanied by a precipitation of a part of the dissolved ones; and when such chemical alternations surpass a definite limit death ensues.—C. R. B.

Sap-flow.—WIEGAND gives a useful résumé of the facts and theories in regard to the flow of the sap from bleeding trees, especially the maple, to which he adds some observations of his own. He holds that the bleeding due to pressure of expanding gases is inadequate to account for the volume of sap exuded, and that the best explanation is the one that ascribes the pressure to the excretion of water by cells of the medullary rays, stimulated to activity by rising temperature.—C. R. B.

Glycogen and paraglycogen.—Since Massart's edition of Errera's post-humous paper on this subject,³² drawings have been found and copied with "scrupulous fidelity." These are now issued as a supplementary paper.³³ The five plates are handsome chromolithographs showing the distribution of glycogen and paraglycogen in the rhizopods and flagellates, as well as several groups of fungi, including the Myxomycetes.—RAYMOND H. POND.

Cotyledons of Trollius.—Ramaley³⁴ has found that the petioles of the cotyledons of *Trollius albiflorus* are united for about half their length, the plumule escaping through a slit in this sheath. Lubbock described a similar condition in *T. Ledebouri*, and it seems to be general throughout Ranunculaceae.—J. M. C.

Mycorhiza and nitrogen fixation.—Contrary to the supposition of MÜLLER, MÖLLER finds that the dichotomous mycorhiza of the mountain pine is of no use in fixing free nitrogen for the tree. 35—RAYMOND H. POND.

Anatomy of roots.—Holm³⁶ has described and illustrated the anatomical structure of the roots of species of Spigelia, Phlox, and Ruellia.—J. M. C.

³º Gorke, H., Ueber chemische Vorgänge beim Erfrieren der Pflanzen. Landw. Versuchs-Stat. **65:**149–160. 1906.

³¹ Wiegand, K. M., Pressure and flow of sap in the maple. Amer. Nat. 40: 409-453. 1906.

³² See review in Bot. GAZETTE 41:370. 1906.

³³ Errera, Leo, Dessins relatifs au glycogen et au paraglycogen. Recueil de l'Institut bot. Bruxelles 1:432–436. pls. 5. 1906.

³⁴ RAMALEY, FRANCIS, The seed and seedling of the mountain globe-flower. Univ. Colorado Studies 3:93-95. figs. 13. 1906.

³⁵ MÖLLER, A., Mykorhizen und Stickstoffernährung. Ber. Deutsch. Bot. Gesells. 24:230–233. 1906.

³⁶ HOLM, THEO., The root-structure of *Spigelia marilandica* L., *Phlox ovata* L., and *Ruellia ciliosa* Pursh. Amer. Jour. Pharmacy **78**:553-559. *figs*. 5. 1906.